

ANNIHILATORS IN BCK-ALGEBRAS II

MICHIRO KONDO

(Received: December 10, 1997)

ABSTRACT. In this paper we show that for every *BCI*-algebra X ,

(1) the family $CI(X)$ of all closed ideal in X forms a pseudo-complemented distributive lattice if and only if X is the *BCK*-algebra;

(2) $\forall A \in CI(X) A \subseteq A^{**} \iff X$ is the *BCK*-algebra ;

1. BCK-ALGEBRA

Firstly we define a *BCI*-algebra and a *BCK*-algebra. An algebra $(X; *, 0)$ of type (2,0) is called a *BCI*-algebra when it satisfies the conditions: For every $x, y, z \in X$,

- (1) $(x * y) * (x * z) \leq z * y$
- (2) $x * (x * y) \leq y$
- (3) $x * x = 0$
- (4) $x * y = y * x = 0 \implies x = y$,

,where the relation " \leq " is defined by

$$x \leq y \text{ if and only if } x * y = 0.$$

It is well known that the relation is a partially ordered relation on any *BCI*-algebra (cf [2]).

In *BCI*-algebras, the following properties hold:

- (a) $x * 0 = x$
- (b) $0 * (x * y) = (0 * x) * (0 * y)$
- (c) $(x * y) * z = (x * z) * y$
- (d) $(x * z) * (y * z) \leq x * y$
- (e) $x \leq y \implies x * z \leq y * z$
- (f) $x \leq y \implies z * x \leq z * y$

We say that a *BCI*-algebra X is a *BCK*-algebra if

- (5) $0 * x = 0$ for $x \in X$.

By an ideal we mean a subset I of X such that it satisfies the conditions:

- (I1) $0 \in I$
- (I2) $x * y, y \in I \implies x \in I$.

1991 *Mathematics Subject Classification.* 06F35, 06D15.

Key words and phrases. annihilator, *BCK*-algebra.

An ideal I is called closed whenever $0 * x \in I$ for any $x \in I$. By $CI(X)$, we denote the set of all closed ideals of X .

For every non-empty subset A of X , we define a subset A^* which is called an annihilator of A :

$$A^* = \{x \in X \mid \forall a \in A (a * (a * x) = 0)\}.$$

The set A is called involutory if $A = A^{**}$. By $S(X)$ we mean the set of all involutory ideals of X . It is proved in [1] that if X is a BCK -algebra and A is a non-empty subset of X then A^* is the ideal of X . We can show the same result for BCI -algebras.

Proposition 1. *Let A be a non-empty subset of a BCI -algebra X . Then A^* is a closed ideal of X .*

Proof. For every $a \in A$, since $a * (a * 0) = a * a = 0$, we have $0 \in A^*$. Next we suppose that $x, y * x \in A^*$. We obtain from definition that

$$\begin{aligned} (6) \quad & a * (a * x) = 0 \text{ and} \\ (7) \quad & a * (a * (y * x)) = 0. \end{aligned}$$

It follows from (2) and (6) that $0 * x = 0$ and hence $(a * x) * a = (a * a) * x = 0 * x = 0$. This means that $a = a * x$ by (4). Similarly we have $a = a * (y * x)$. From these, we have in turn

$$\begin{aligned} a &= a * (y * x) = (a * x) * (y * x) \leq a * y \quad (\text{by } (d)) \\ 0 &= a * a \leq (a * y) * a = 0 * y \quad (\text{by } (e)) \\ 0 &= 0 * (0 * y) \leq y, \text{ which implies that } 0 * y = 0. \end{aligned}$$

It follows that $(a * y) * a = (a * a) * y = 0 * y = 0$ and hence $a = a * y$. Thus $y \in A^*$ and A^* is the ideal of X .

Lastly we show that A^* is closed, that is, $0 * x \in A^*$ whenever $x \in A^*$. For $x \in A^*$, we obtain that $0 = a * (a * x) \leq x$ for every $a \in A$. Thus we have $0 * x = 0 \in A^*$. This means that A^* is closed.

Let X be a BCI -algebra. We define a subset G called BCK -part of X as

$$G = \{x \in X \mid 0 * x = 0\}$$

For the subset G it is proved that

Proposition 2. (1) G is the closed ideal, that is, $G \in CI(X)$.

$$(2) \quad G = \{0\}^*$$

$$(3) \quad G^* = \{0\}$$

Proof. We only prove the case of (1). The other cases are proved similarly (cf. [3]). It is obvious that $0 \in G$. If $x, y * x \in G$, then $0 * x = 0 * (y * x) = 0$. Since $0 * (y * x) = (0 * y) * (0 * x)$, we have $0 * y = 0$. Thus G is the ideal. Moreover, for any $x \in G$, it follows from definition of G that $0 * x = 0 \in G$. This means that G is closed.

Proposition 3. *Let X be a BCI -algebra and A a subset of X . Then we have*

$$0 \in A \text{ if and only if } A \cap A^* = \{0\}$$

Proof. We show the "only if" part. We suppose that $0 \in A$. First of all we note that $A \cap A^* \neq \emptyset$ because of $0 \in A^*$. For every $x \in A \cap A^*$, we have

$x * (x * x) = 0$ by definition of A^* . On the other hand $x * (x * x) = x * 0 = x$. It follows that $x = 0$ and hence $A \cap A^* = \{0\}$.

Corollary 1. $A \cap A^* = \emptyset$ or $A \cap A^* = \{0\}$

Hence we have $A \cap A^* = \emptyset$ for every ideal A in any BCI -algebra (cf [1]).

2. STRUCTURE OF $I(X)$

In this section we consider the set $I(X)$ of all ideals of a BCK -algebra X . In case of the BCK -algebra X , every ideal of X is identical with the closed ideal, that is, $I(X) = CI(X)$. In [4], it is proved that if X is a BCK -algebra then the set $I(X)$ of all ideals of X forms the pseudo-complemented distributive lattice. In this section we show the converse, that is, for every BCI -algebra X , if $CI(X)$ is the pseudo-complemented distributive lattice then X is the BCK -algebra.

Proposition 4. $A \subseteq B \implies B^* \subseteq A^*$

Proof. [3]

We consider the following condition (BCK) which is proved to be equivalent to that X is a BCK -algebra ([4]):

(BCK) For every ideal A , if $x \in X$ and $a \in A$ then $a * (a * x) \in A$.

Proposition 5. Let X be any BCI -algebra. Then we have
 $X : BCK\text{-algebra} \iff \text{the condition (BCK) holds.}$

Lemma 1. Let X be a BCK -algebra and $A, B \in I(X)$.

$A \cap B = \{0\} \iff A \subseteq B^*$.

Proof. See [4]

Lemma 2. If X is a BCK -algebra, then $A \subseteq A^{**}$ for every $A \in I(X)$.

Proof. Let $a \in A$. For any $x \in A^*$, since A^* is the ideal, it follows from (BCK) that $x * (x * a) \in A^*$. On the other hand, since $x * (x * a) \leq a \in A$, we have $x * (x * a) \in A \cap A^* = \{0\}$. This implies that $x * (x * a) = 0$ for every $x \in A^*$. Hence we have $A \subseteq A^{**}$.

Remark: In [1] the results above are proved for every commutative BCK -algebra. But we see that the commutativity is unnecessary.

Lemma 3. Let X be a BCI -algebra. Then

$\{0\}^* = X \iff X : BCK\text{-algebra.}$

Proof. For any $x \in X = \{0\}^*$, we have $0 * (0 * x) = 0$ by definition of annihilator. This means that $0 * x = 0$, that is, X is the BCK -algebra.

The converse is trivial.

Theorem 1. Let X be a BCI -algebra. Then

$\forall A \in CI(X) (A \subseteq A^{**}) \iff X : BCK\text{-algebra}$

Proof. Suppose that $A \subseteq A^{**}$ for every $A \in CI(X)$. Clearly X is a closed ideal of X , we have $X \subseteq X^{**}$ and hence $X = X^{**} = \{0\}^*$. It follows from lemma 3 that X is the BCK -algebra.

In [6], it is proved that the set $I(X)$ of all ideals of BCK -algebra X is the distributive lattice, where

$$\begin{aligned} A \wedge B &= A \cap B \\ A \vee B &= \{x \in X \mid \exists a \in A, \exists b \in B; (x * a) * b = 0\}. \end{aligned}$$

It is clear from the argument above that the set $I(X) = CI(X)$ of all ideals of BCK -algebra X forms the pseudo-complemented distributive lattice (cf [4]). In the following we show that the converse holds, that is, if $CI(X)$ is a pseudo-complemented distributive lattice then X is the BCK -algebra. We also give an example which indicates that the pseudo-complementedness of $CI(X)$ is essential.

Theorem 2. *Let X be a BCI -algebra. If $(CI(X); \wedge, \vee, \phi, X)$ is the pseudo-complemented distributive lattice, then X is the BCK -algebra.*

Proof. We assume that $CI(X)$ is the pseudo-complemented distributive lattice. For every closed ideal $A \in CI(X)$, we have

$$\begin{aligned} A \wedge (A \vee A^{**}) &= (A^* \wedge A) \vee (A^* \wedge A^{**}) \\ &= \{0\} \vee \{0\} \\ &= \{0\}. \end{aligned}$$

By pseudo-complementedness, it follows that $A \vee A^{**} \subseteq A^{**}$ and hence $A \vee A^{**} = A^{**}$. This implies that $A \subseteq A^{**}$ for every $A \in CI(X)$. Thus X is the BCK -algebra.

By the example below we know that the pseudo-complementedness is essential, that is, X is not need the BCK -algebra even if $CI(X)$ is distributive.

Example

*	0	1	2	3
0	0	0	3	3
1	1	0	3	2
2	2	3	0	1
3	3	3	0	0

It is easy to see that $CI(X) = \{\{0\}, \{0, 1\}, \{0, 3\}, X\}$ and hence that $CI(X)$ is a distributive lattice. The algebra is not a BCK -algebra, because we have $\{0, 1\} \cap \{0, 3\} = \{0\}$ but $\{0, 3\}$ is not a subset of $\{0, 1\}^* = \{0\}$.

It is proved in [4] that the set $S(X) = (S(X); \wedge, \sqcup, *, \phi, X)$ of all involutory ideals of a BCK -algebra X forms a Boolean algebra, where $A \sqcup B$ is defined as $(A^* \cap B^*)^*$ for every $A, B \in S(X)$. If we consider the map $\xi : I(X) \rightarrow S(X)$ defined by $\xi(A) = A^*$, then we have $\ker \xi$ is a congruence over $I(X)$. From the general theory of universal algebras, we can conclude that

Theorem 3. *$I(X)/\ker \xi \cong S(X)$. Hence $I(X)/\ker \xi$ is the Boolean algebra.*

REFERENCES

- [1] M. Aslam and A. B. Thaheem, *On certain ideals in BCK-algebras*, Math. Japonica vol. **36** (1991), 895-906.
- [2] K. Iseki and S. Tanaka, *An introduction to the theory of BCK-algebras*, Math. Japonica vol. **23** (1978), 1-26.
- [3] Y. B. Jun, E. H. Roh, and J. Meng, *Annihilators in BCI-algebras*, Math. Japonica vol. **43** (1996), 559-562.
- [4] M. Kondo, *Annihilators in BCK-algebras*, submitted.
- [5] M. Palasinski, *On ideals and congruence lattices of BCK-algebras*, Math. Seminar Notes vol. **9** (1981), 441-443.
- [6] S. M. Wei and Y. B. Jun, *Ideal lattices of BCI-algebras*, Math. Japonica vol. **44** (1996), 303-305.

DEPARTMENT OF MATHEMATICS, SHIMANE UNIVERSITY, MATSUE 690-8504 JAPAN